




Global bifurcation and nodal solutions for homogeneous Kirchhoff type equations

Fang Liu¹, Hua Luo² and Guowei Dai ¹

¹School of Mathematical Sciences, Dalian University of Technology, Dalian, 116024, PR China

²School of Economics and Finance, Shanghai International Studies University, Shanghai, 201620, PR China

Received 20 December 2019, appeared 6 May 2020

Communicated by Maria Alessandra Ragusa

Abstract. In this paper, we shall study unilateral global bifurcation phenomenon for the following homogeneous Kirchhoff type problem

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right) u'' = \lambda u^3 + h(x, u, \lambda) & \text{in } (0, 1), \\ u(0) = u(1) = 0. \end{cases}$$

As application of bifurcation result, we shall determine the interval of λ in which there exist nodal solutions for the following homogeneous Kirchhoff type problem

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right) u'' = \lambda f(x, u) & \text{in } (0, 1), \\ u(0) = u(1) = 0, \end{cases}$$

where f is asymptotically cubic at zero and infinity. To do this, we also establish a complete characterization of the spectrum of a homogeneous nonlocal eigenvalue problem.

Keywords: bifurcation, spectrum, nonlocal problem, nodal solution, regularity results.

2020 Mathematics Subject Classification: 34C23, 47J10, 34C10.

1 Introduction

Consider the following problem

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right) u'' = \lambda u^3 + h(x, u, \lambda) & \text{in } (0, 1), \\ u(0) = u(1) = 0, \end{cases} \quad (1.1)$$

where λ is a nonnegative parameter and $h : (0, 1) \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function satisfying

$$\lim_{s \rightarrow 0} \frac{h(x, s, \lambda)}{s^3} = 0 \quad (1.2)$$

 Corresponding author. Email: daiguowei@dlut.edu.cn

uniformly for all $x \in (0, 1)$ and λ on bounded sets.

The problem (1.1) is related to the stationary problem of a model introduced by Kirchhoff in 1883 to describe the transversal oscillations of a stretched string [16]. Some important and interesting results can be found, for example, in [1, 4, 12, 13, 15, 19, 25]. Recently, there are many mathematicians studying the problem (1.1), see [5, 6, 8, 17, 20, 21, 22, 24, 26] and the references therein. A distinguishing feature of problem (1.1) is that the first equation contains a nonlocal coefficient $\int_0^1 |u'|^2 dx$, and hence the equation is no longer a pointwise identity, which raises some essential difficulties to the study of this kind of problems. In particular, the bifurcation theory of [11, 23] does not work on it.

As shown in [3], the following problem

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right) u'' = \lambda u^3 & \text{in } (0, 1), \\ u(0) = u(1) = 0 \end{cases} \quad (1.3)$$

possesses infinitely many eigenvalues $0 < \mu_1 < \mu_2 < \dots < \mu_k \rightarrow +\infty$, all of which are simple. The eigenfunction φ_k corresponding to μ_k has exactly $k - 1$ simple zeros in $(0, 1)$. Let S_k^+ denote the set of functions in $E := C_0^1[0, 1]$ which have exactly $k - 1$ interior nodal (i.e. non-degenerate) zeros in $(0, 1)$ and are positive near $x = 0$, and set $S_k^- = -S_k^+$, and $S_k = S_k^+ \cup S_k^-$. It is clear that S_k^+ and S_k^- are disjoint and open in E . Finally, let $\Phi_k^\pm = \mathbb{R} \times S_k^\pm$ and $\Phi_k = \mathbb{R} \times S_k$ under the product topology. The first main result of this paper is the following theorem.

Theorem 1.1. *The pair $(\mu_k, 0)$ is a bifurcation point of (1.1). Moreover, there are two distinct unbounded continua in $\mathbb{R} \times H_0^1(0, 1)$, \mathcal{C}_k^+ and \mathcal{C}_k^- , consisting of the bifurcation branch \mathcal{C}_k emanating from $(\mu_k, 0)$, such that $\mathcal{C}_k^\nu \subseteq (\{(\mu_k, 0)\} \cup \Phi_k^\nu)$, $\nu \in \{+, -\}$.*

It is well known that the index formula of an isolated zero is very important in the study of bifurcation phenomena for semi-linear differential equations. However, problem (1.1) is nonlinear. In order to overcome this difficulty, we study the following auxiliary homogeneous eigenvalue problem

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right)^{p/2} u'' = \lambda |u|^p u & \text{in } (0, 1), \\ u(0) = u(1) = 0, \end{cases} \quad (1.4)$$

where $p \in [0, 2]$. We study the spectral structure, and establish an index formula via a suitable homotopic deformation from a general $p \in [0, 2]$ to $p = 0$ for problem (1.4). Let $\lambda_1(p)$ denote the first eigenvalue of (1.4). As shown in [9], $\lambda_1(p) > 0$ is simple, isolated, the unique principal eigenvalue of (1.4), and is continuous with respect to p . Our second main result is the following theorem.

Theorem 1.2. *The set of all eigenvalues of (1.4) is formed by a sequence*

$$0 < \lambda_1(p) < \lambda_2(p) < \dots < \lambda_k(p) \rightarrow +\infty.$$

Every $\lambda_k(p)$ is simple, continuous with respect to p and the corresponding one dimensional space of solutions of (1.4) with $\lambda = \lambda_k(p)$ is spanned by a function having precisely k bumps in $(0, 1)$. Each k -bump solution is constructed by the reflection and compression of the eigenfunction φ_1 associated with $\lambda_1(p)$.

Based on Theorem 1.1, we study the existence of nodal solutions for the following problem

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right) u'' = \lambda f(x, u) & \text{in } (0, 1), \\ u(0) = u(1) = 0. \end{cases} \quad (1.5)$$

We assume that f satisfies the following conditions

(f1) $f : (0, 1) \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function such that $f(x, s)s > 0$ for all $x \in (0, 1)$ and any $s \neq 0$.

(f2) there exist $f_0, f_\infty \in (0, +\infty)$ such that

$$\lim_{s \rightarrow 0^+} \frac{f(x, s)}{s^3} = f_0, \quad \lim_{s \rightarrow +\infty} \frac{f(x, s)}{s^3} = f_\infty$$

uniformly with respect to all $x \in (0, 1)$.

The last main theorem of this paper is the following result.

Theorem 1.3. *Assume that f satisfies (f1)–(f2). Then the pair $(\mu_k/f_0, 0)$ is a bifurcation point of (1.5) and there are two distinct unbounded continua in $\mathbb{R} \times H_0^1(0, 1)$, \mathcal{C}_k^+ and \mathcal{C}_k^- , emanating from $(\mu_k/f_0, 0)$, such that $\mathcal{C}_k^\nu \subseteq (\{(\mu_k/f_0, 0)\} \cup \Phi_k^\nu)$ and links $(\mu_k/f_0, 0)$ to $(\mu_k/f_\infty, \infty)$.*

The rest of this paper is arranged as follows. In Section 2, we establish the spectrum of problem (1.4). In Section 3 and 4, we give the proofs of Theorem 1.1 and 1.3, respectively.

2 Spectrum of (1.4)

Let X be the usual Sobolev space $H_0^1(0, 1)$ with the norm $\|u\| = (\int_0^1 |u'|^2 dx)^{1/2}$. For any $\alpha \in (0, 1]$, we use $C^\alpha[0, 1]$ to denote all the real functions such that

$$\|u\|_\alpha := \sup_{x, y \in [0, 1], x \neq y} \frac{|u(x) - u(y)|}{|x - y|^\alpha} < +\infty.$$

Firstly, we have the following regularity result.

Proposition 2.1. *Any weak solution $u \in X$ of problem (1.4) is also a classical solution, i.e., $u \in C^2[0, 1]$ satisfying (1.4).*

Proof. Let u be a nontrivial weak solution of problem (1.4) and

$$f(x) = \frac{\lambda |u(x)|^p u(x)}{\|u\|^p}.$$

Note that

$$H_0^1(0, 1) = \{u \in AC[0, 1] : u' \in L^2(0, 1) \text{ and } u(0) = u(1) = 0\}.$$

Then it is obvious that $f \in L^2(0, 1)$, in fact continuous by the compact embedding $X \hookrightarrow C^{1/2}[0, 1]$. According to the definition of weak solution, we have

$$-\left(\int_0^1 |u'|^2 dx\right)^{\frac{p}{2}} u'' = \lambda |u|^p u$$

in the sense of distribution. It follows that

$$u'(x) = u'(0) - \int_0^x f(t) dt.$$

Note that

$$u(x) = \int_0^x u'(t) dt.$$

So, we have that

$$u(x) = \int_0^x \left(u'(0) - \int_0^t f(\tau) d\tau \right) dt = u'(0)x - \int_0^x \int_0^t f(\tau) d\tau dt.$$

Then, in view of $f \in C[0, 1]$, we get that $u \in C^2[0, 1]$ and satisfies (1.4). \square

Lemma 2.2. *If (λ, u) is a solution of (1.4) and u has a double zero, then $u \equiv 0$.*

Proof. Let u be a solution of (1.4) and $x^* \in [0, 1]$ be a double zero. If $\|u\| = 0$, the conclusion is obvious. Next, we assume that $\|u\| \neq 0$. We note that

$$u(x) = -\frac{\lambda}{\|u\|^p} \int_{x^*}^x \int_{x^*}^s |u|^p u d\tau ds.$$

Firstly, we consider $x \in [0, x^*]$. Then

$$\begin{aligned} |u(x)| &= \left| -\frac{\lambda}{\|u\|^p} \int_{x^*}^x \int_{x^*}^s |u|^p u d\tau ds \right| \leq \left| \frac{\lambda}{\|u\|^p} \int_{x^*}^x \int_{x^*}^x |u|^p u d\tau ds \right| \\ &= \left| \frac{\lambda}{\|u\|^p} (x - x^*) \int_{x^*}^x |u|^p u d\tau \right| \\ &\leq \frac{\lambda}{\|u\|^p} \int_x^{x^*} |u|^{p+1} d\tau \leq \frac{\lambda \|u\|_\infty^p}{\|u\|^p} \int_x^{x^*} |u| d\tau \leq \lambda \int_x^{x^*} |u| d\tau. \end{aligned}$$

By the Gronwall–Bellman inequality [7, Lemma 2.2], we get $u \equiv 0$ on $[0, x^*]$. Similarly, we can get $u \equiv 0$ on $[x^*, 1]$ and the proof is completed. \square

Lemma 2.3. *Each nontrivial solution (λ, u) of (1.4) has a finite number of zeros.*

Proof. Suppose, on the contrary, that u has a sequence zeros x_n . Since $[0, 1]$ is compact, up to a subsequence, there exists $x_0 \in [0, 1]$ such that $\lim_{n \rightarrow +\infty} x_n = x_0$. By the continuity of u , we have that $u(x_0) = \lim_{n \rightarrow +\infty} u(x_n) = 0$. So, we have that

$$u'(x_0) = \lim_{n \rightarrow +\infty} \frac{u(x_n) - u(x_0)}{x_n - x_0} = 0.$$

Thus, x_0 is a double zero of u . By Lemma 2.2, we get that $u \equiv 0$, which is a contradiction. \square

Let J be a strict sub-interval of I . Let $\lambda_1(J)$ denote the first eigenvalue

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right)^{p/2} u'' = \lambda |u|^p u & \text{in } J, \\ u(x) = 0 & \text{on } \partial J, \end{cases}$$

where $p \in [0, 2]$.

Lemma 2.4. $\lambda_1(I)$ verifies the strict monotonicity property with respect to the domain I , i.e. if J is a strict subinterval of I , then $\lambda_1(I) < \lambda_1(J)$.

Proof. Let φ_1 with $\|\varphi_1\| = 1$ be the eigenfunction of (1.4) on J corresponding to $\lambda_1(J)$, and denote by $\tilde{\varphi}_1$ the extension by zero on I . Then we have that

$$\frac{1}{\lambda_1(J)} = \int_J |\varphi_1|^{p+2} dx = \int_I |\tilde{\varphi}_1|^{p+2} dx < \sup_{u \in X, \|u\|=1} \int_0^1 |u|^{p+2} dx = \frac{1}{\lambda_1(I)}.$$

The last strict inequality holds from the fact that $\tilde{\varphi}_1$ vanishes in $I \setminus J$ so cannot be an eigenfunction corresponding to the principal eigenvalue $\lambda_1(I)$. \square

Proof of Theorem 1.2. Let φ_1 be a positive eigenfunction corresponding to $\lambda_1(p)$. It follows from the symmetry of (1.4) and Theorem 3.1 of [9] (or Theorem 2.4 of [18]) that $\varphi_1(x) = \varphi_1(1-x)$ for $x \in [0, 1]$, i.e. φ_1 is even with respect to $1/2$. For any $k \geq 2$, set

$$\varphi_k(x) = \begin{cases} \varphi_1(kx), & x \in [0, \frac{1}{k}], \\ -\varphi_1(kx-1), & x \in [\frac{1}{k}, \frac{2}{k}], \\ \vdots & \vdots \\ (-1)^k \varphi_1(kx-k+1), & x \in [\frac{k-1}{k}, 1]. \end{cases}$$

Then φ_k is an eigenfunction of (1.4) associated with the eigenvalue $\lambda_k(p) = k^{p+2} \lambda_1(p)$. Clearly, the continuity of $\lambda_1(p)$ implies that $\lambda_k(p)$ is continuous with respect to p .

On the other hand, let $u = u(x)$ be an eigenfunction of (1.4) associated with some eigenvalue $\lambda_* > \lambda_1(p)$. According to Theorem 3.1 of [9], u changes sign in $(0, 1)$. Lemmas 2.2 and 2.3 imply that $u \in S_k$ for some $k \geq 2$. Without loss of generality, we may assume that $u'(0) > 0$. Let

$$0 < \tau_1 < \tau_2 < \cdots < \tau_{k-1} < 1$$

denote the zeros of u in $(0, 1)$. Without loss of generality, we may assume that $\tau_1 \leq 1/k$. Applying Lemma 2.4 on $[0, 1/k]$, we have that $\lambda_* \geq \lambda_k$. By Lemma 2 of [2], there exist integers p and q , $1 \leq p \leq k-1$, $1 \leq q \leq k-1$, such that

$$\tau_p \leq \frac{1}{q+1} < \frac{1}{q} \leq \tau_{p+1}.$$

Applying Lemma 2.4 on $[\tau_p, \tau_{p+1}]$, we have that $\lambda_* \leq \lambda_k$. So we have that $\lambda_* = \lambda_k$. Furthermore, if $\tau_1 < 1/k$, we have that $\lambda_* > \lambda_k$; if $\tau_1 > 1/k$, we have that $\lambda_* < \lambda_k$. Thus we have $\tau_1 = 1/k$ and $u = c_1 \varphi_k(x)$ for $x \in [0, 1/k]$. Similarly, we can obtain that $\tau_i = i/k$ and $u = c_i \varphi_k(x)$ for $x \in [(i-1)/k, i/k]$, $2 \leq i \leq k-1$. Let us normalize u as $u'(0) = \varphi'_k(0)$. It follows that $c_1 = 1$. Hence $\varphi'_k(\frac{1}{k}) = c_2 \varphi'_k(\frac{1}{k})$. So we have $c_2 = 1$. Similarly, one has $c_i = 1$ for all $3 \leq i \leq k-1$. Therefore, we have that $u(x) = \varphi_k(x)$, $x \in [0, 1]$. \square

3 Global bifurcation

Consider the following auxiliary problem

$$\begin{cases} - \left(\int_0^1 |u'|^2 dx \right)^{p/2} u'' = f(x) & \text{in } (0, 1), \\ u(0) = u(1) = 0 \end{cases} \quad (3.1)$$

for any $p \in [0, 2]$ and a given $f \in X^*$. We have shown in [9] that problem (3.1) has a unique weak solution. Let us denote by $R_p(f)$ the unique weak solution of (3.1). Then $R_p : X^* \rightarrow X$

is a continuous operator. Since the embedding of $X \hookrightarrow L^\infty(0,1)$ is compact, the restriction of R_p to $L^1(0,1)$ is a completely continuous (i.e., continuous and compact) operator. From the obvious modification of Lemma 4.2 of [9], we can get the following compactness and continuity of the operator R_p with respect to p and f .

Lemma 3.1. *The operator $R : [0,2] \times L^1(0,1) \rightarrow L^\infty(0,1)$ defined by $R(p,f) = R_p(f)$ is completely continuous.*

Now, we consider (1.4) again. Clearly, u is a weak solution of (1.4) if and only if $u \in X$, $\lambda \in [0, +\infty)$ satisfy

$$u = R_p(\lambda|u|^p u) = \lambda^{\frac{1}{p+1}} R_p(|u|^p u) := T_p^\lambda(u).$$

For any $u \in X$, we define

$$K_p(u) = |u|^p u.$$

Then we see that $K_p(u) \in L^1(0,1)$. We claim that $K_p : X \hookrightarrow L^1(0,1)$ is continuous. Assume that $u_n \rightarrow u$ in X . Since embedding $X \hookrightarrow C[0,1]$ is compact, we have $u_n \rightarrow u$ in $C[0,1]$. It follows that $u_n(x) \rightarrow u(x)$ for any $x \in [0,1]$. So, we have that $K_p(u_n) \rightarrow K_p(u)$ in $L^1(0,1)$. Since $R_p : L^1(0,1) \rightarrow X$ is a compact, we have that $T_p^\lambda = \lambda^{\frac{1}{p+1}} R_p \circ K_p : X \rightarrow X$ is completely continuous. Thus the Leray–Schauder degree

$$\deg_X(I - T_p^\lambda, B_r(0), 0)$$

is well-defined for arbitrary r -ball $B_r(0)$ and $\lambda \neq \lambda_k(p)$. It is well known that

$$\deg_X(I - T_0^\lambda, B_r(0), 0) = (-1)^\beta,$$

where β is the number of eigenvalues of problem (1.4) with $p = 0$ less than λ . As far as the general p , we can compute it through the deformation along p .

Proposition 3.2. *Let $r > 0$ and $\bar{p} \in [0,2]$. Then*

$$\deg_X(I - T_{\bar{p}}^\lambda, B_r(0), 0) = \begin{cases} 1, & \text{if } \lambda \in (0, \lambda_1(\bar{p})), \\ (-1)^k, & \text{if } \lambda \in (\lambda_k(\bar{p}), \lambda_{k+1}(\bar{p})). \end{cases}$$

Proof. If $\lambda \in (0, \lambda_1(\bar{p}))$, the conclusion has done in [9]. So we only need to prove the case $\lambda \in (\lambda_k(\bar{p}), \lambda_{k+1}(\bar{p}))$. Since $p \rightarrow \lambda_k(p)$ is continuous, we can define a continuous function $\chi : [0,2] \rightarrow \mathbb{R}$ such that $\lambda_k(p) < \chi(p) < \lambda_{k+1}(p)$ and $\lambda = \chi(\bar{p})$. Set

$$d(p) = \deg_X(I - T_p^{\chi(p)}, B_r(0), 0).$$

We shall show that $d(p)$ is constant in $[0,2]$.

Define $S_p : L^\infty(0,1) \rightarrow X$ by $S_p(u) = R_p(\chi(p)|u|^p u)$. We see that $S_p(u) = \chi^{\frac{1}{p+1}}(p) R_p \circ K_p(u)$, where $K_p(u) = |u|^p u$. By the definition of K_p , we can easily verify that $K_p : L^\infty(0,1) \rightarrow L^1(0,1)$ is continuous. Since $R_p : L^1(0,1) \rightarrow X$ is a compact, we get that $S_p : L^\infty(0,1) \rightarrow X$ is completely continuous. Also we have that $T_p^{\chi(p)} = S_p \circ i$ where $i : X \rightarrow L^\infty(0,1)$ is the usual inclusion. From Lemma 2.4 of [14], we obtain that

$$d(p) = \deg_{L^\infty}(I - i \circ S_p, \Omega_s, 0) \quad \text{for } p \in [0,2],$$

where Ω_s is any open bounded set in $L^\infty(0,1)$ containing 0. It is not difficult to verify that the operator $\varphi : [0,2] \times L^\infty(0,1) \rightarrow L^1(0,1)$ defined by $\varphi(p, u) = |u|^p u$ is continuous. This fact, the continuity of $\chi(p)$ and Lemma 3.1 imply that $(p, u) \mapsto R_p(\chi(p)|u|^p u) = (i \circ S_p)(u) : [0,2] \times L^\infty(0,1) \rightarrow L^\infty(0,1)$ is completely continuous. Since $\lambda_k(p) < \chi(p) < \lambda_{k+1}(p)$ for any $p \in [0,2]$, we have that $u - R_p(\chi(p)|u|^p u) \neq 0$ on $\partial\Omega_s$. The invariance of the Leray–Schauder degree under a compact homotopy follows that $d(p) \equiv \text{constant}$ for $p \in [0,2]$. So, $d(\bar{p}) = d(0) = (-1)^k$, as desired. \square

In particular, we have the following corollary.

Corollary 3.3. *Let $r > 0$. Then*

$$\deg_X \left(I - T_2^\lambda, B_r(0), 0 \right) = \begin{cases} 1, & \text{if } \lambda \in (0, \mu_1), \\ (-1)^k, & \text{if } \lambda \in (\mu_k, \mu_{k+1}), \end{cases}$$

where μ_k is the k -th eigenvalue of (1.3).

Clearly, the pair (λ, u) is a solution of (1.1) if and only if (λ, u) satisfies

$$u = R_2(\lambda u^3 + h(x, u, \lambda)) := G_\lambda(u).$$

It is easy to see that $G_\lambda : X \rightarrow X$ is completely continuous and $G_\lambda(0) = 0, \forall \lambda \in [0, +\infty)$. μ_k is the λ_k . Let X_0 be any complement of $\text{span}\{\varphi_k\}$ in X .

Theorem 3.4. *The pair $(\mu_k, 0)$ is a bifurcation point of (1.1). Moreover, there are two distinct continua in $\mathbb{R} \times X$, \mathcal{C}_k^+ and \mathcal{C}_k^- , consisting of the bifurcation branch \mathcal{C}_k emanating from $(\mu_k, 0)$, which contain $\{(\mu_k, 0)\}$ and each of them satisfies one of the following non-excluding alternatives:*

1. *it is unbounded in $\mathbb{R} \times X$;*
2. *it contains a pair $(\mu_j, 0)$ with $j \neq k$;*
3. *it contains a point $(\lambda, y) \in \mathbb{R} \times (X_0 \setminus \{0\})$.*

Proof. We use the abstract bifurcation result of [10] to prove this theorem. An operator L defined on X is called homogeneous if $L(cu) = cL(u)$ for any $c \in \mathbb{R}$ and $u \in X$. It is not difficult to verify that $L(\lambda) := T_2^\lambda : X \rightarrow X$ is homogeneous and completely continuous. Let $\tilde{h}(x, u, \lambda) = \max_{0 \leq |s| \leq u} |h(x, s, \lambda)|$ for all $x \in (0, 1)$ and λ on bounded sets, then \tilde{h} is nondecreasing with respect to u and

$$\lim_{u \rightarrow 0^+} \frac{\tilde{h}(x, u, \lambda)}{u^3} = 0. \quad (3.2)$$

Further it follows from (3.2) that

$$\frac{h(x, u, \lambda)}{\|u\|^3} \leq \frac{\tilde{h}(x, |u|, \lambda)}{\|u\|_\infty^3} \leq \frac{\tilde{h}(x, \|u\|_\infty, \lambda)}{\|u\|_\infty^3} \rightarrow 0 \quad \text{as } \|u\| \rightarrow 0 \quad (3.3)$$

uniformly for $x \in (0, 1)$ and λ on bounded sets. Let

$$H(\lambda, u) = G_\lambda(u) - L(\lambda)u.$$

By (3.3), we can easily verify that $H : \mathbb{R} \times X \rightarrow X$ is completely continuous with $H = o(\|u\|)$ near $u = 0$ uniformly on bounded λ intervals. Noting Corollary 3.3, the desired conclusions can be obtained by applying Theorem 1 of [10]. \square

By an argument similar to that of Proposition 2.1, we can get the following regularity result.

Proposition 3.5. *Any weak solution $u \in X$ of problem (1.1) is also a classical solution, i.e., $u \in C^2(0, 1) \cap C^{1,\alpha}[0, 1]$ satisfying (1.1) and $u(0) = u(1) = 0$.*

Lemma 3.6. *If (λ, u) is a solution of (1.1) and u has a double zero, then $u \equiv 0$.*

Proof. Let u be a solution of (1.1) and $x^* \in [0, 1]$ be a double zero. If $\|u\| = 0$, the conclusion is done. Next, we assume that $\|u\| \neq 0$. We note that

$$u(x) = \frac{-1}{\|u\|^2} \int_{x^*}^x \int_{x^*}^s (\lambda u^3 + h(x, u, \lambda)) \, d\tau \, ds.$$

Firstly, we consider $x \in [0, x^*]$. Then

$$\begin{aligned} |u(x)| &\leq \frac{1}{\|u\|^2} \int_x^{x^*} |\lambda u^3 + h(x, u, \lambda)| \, d\tau, \\ &\leq \frac{\|u\|_\infty^2}{\|u\|^2} \int_x^{x^*} \left(|\lambda| + \left| \frac{h(\tau, u(\tau), \lambda)}{u(\tau)} \right| \right) |u(\tau)| \, d\tau. \end{aligned}$$

In view of (1.2), for any $\varepsilon > 0$, there exists a constant $\delta > 0$ such that

$$|h(x, s, \lambda)| \leq \varepsilon |s|$$

uniformly with respect to all $x \in (0, 1)$ and fixed λ when $|s| \in [0, \delta]$. Hence,

$$|u(x)| \leq \int_x^{x^*} \left(|\lambda| + \varepsilon + \max_{s \in [\delta, \|u\|_\infty]} \left| \frac{h(\tau, s, \lambda)}{s^3} \right| \right) |u(\tau)| \, d\tau.$$

By the Gronwall–Bellman inequality [7], we get $u \equiv 0$ on $[0, x^*]$. Similarly, we can get $u \equiv 0$ on $[x^*, 1]$ and the proof is complete. \square

Proof of Theorem 1.1. Lemma 3.1 of [10] implies that there exists a bounded open neighborhood \mathcal{O}_k of $(\mu_k, 0)$ such that $(\mathcal{C}_k^v \cap \mathcal{O}_k) \subseteq (\Phi_k^v \cup \{(\mu_k, 0)\})$ or $(\mathcal{C}_k^v \cap \mathcal{O}_k) \subseteq (\Phi_k^{-v} \cup \{(\mu_k, 0)\})$. Without loss of generality, we assume that $(\mathcal{C}_k^v \cap \mathcal{O}_k) \subseteq (\Phi_k^v \cup \{(\mu_k, 0)\})$.

Next, we show that $\mathcal{C}_k^v \subseteq (\Phi_k^v \cup \{(\mu_k, 0)\})$. Suppose $\mathcal{C}_k^v \not\subseteq (\Phi_k^v \cup \{(\mu_k, 0)\})$. Then there exists $(\mu, u) \in \mathcal{C}_k^v \cap (\mathbb{R} \times \partial S_k^v)$ such that $(\mu, u) \neq (\mu_k, 0)$ and $(\lambda_n, u_n) \rightarrow (\mu, u)$ with $(\lambda_n, u_n) \in \mathcal{C}_k^v \cap (\mathbb{R} \times S_k^v)$. Since $u \in \partial S_k^v$, by Lemma 3.6, $u \equiv 0$. Let $v_n := u_n / \|u_n\|$, then v_n should be a solution of the following problem

$$v = R_2 \left(\lambda_n v^3 + \frac{h(x, u_n, \lambda_n)}{\|u_n(x)\|^3} \right). \quad (3.4)$$

By (3.3), (3.4) and the compactness of R_2 we obtain that for some convenient subsequence $v_n \rightarrow v_0 \neq 0$ as $n \rightarrow +\infty$. Now v_0 verifies the equation

$$-\int_0^1 |v'|^2 \, dx v'' = \mu v^3$$

and $\|v_0\| = 1$. Hence $\mu = \mu_j$, for some $j \neq k$. Hence $v_0 \in S_j$ which is an open set in X , and as a consequence for some n large enough, $u_n \in S_j$, and this is a contradiction. Thus, we have that

$$\mathcal{C}_k^v \subseteq (\Phi_k^v \cup \{(\mu_k, 0)\}).$$

Furthermore, by an argument similar to the above, we can easily show that $\mathcal{C}_k \cap (\mathbb{R} \times \{0\}) = \{(\mu_k, 0)\}$. So Theorem 1 of [10] implies that \mathcal{C}_k is unbounded.

We claim that both \mathcal{C}_k^+ and \mathcal{C}_k^- are unbounded. Introduce the following auxiliary problem

$$\begin{cases} -\left(\int_0^1 |u'|^2 dx\right) u'' = \lambda u^3 + \tilde{h}(x, u, \lambda) & \text{in } (0, 1), \\ u(0) = u(1) = 0, \end{cases}$$

where \tilde{h} is defined by

$$\tilde{h}(x, u, \lambda) = \begin{cases} h(x, u, \lambda), & \text{if } u'(0) > 0, \\ -h(x, -u, \lambda), & \text{if } u'(0) < 0. \end{cases}$$

The previous argument shows that an unbounded continuum $\tilde{\mathcal{C}}_k$ bifurcates from $(\mu_k, 0)$ and can be split into $\tilde{\mathcal{C}}_k^+$ and $\tilde{\mathcal{C}}_k^-$ with $\tilde{\mathcal{C}}_k^v$ connected, $\tilde{\mathcal{C}}_k^v \subseteq (\{(\mu_k, 0)\} \cup (\mathbb{R} \times S_k^v))$. It is easy to see that $\tilde{\mathcal{C}}_k^- = -\tilde{\mathcal{C}}_k^+$. It follows that both $\tilde{\mathcal{C}}_k^+$ and $\tilde{\mathcal{C}}_k^-$ are unbounded. It is clear that $\tilde{\mathcal{C}}_k^+ \subseteq \mathcal{C}_k^+$. Therefore \mathcal{C}_k^+ must be unbounded. A symmetric argument shows that \mathcal{C}_k^- is also unbounded. \square

4 Nodal solutions

In this section, we apply Theorem 1.1 to study the existence of nodal solutions for (1.5).

Proof of Theorem 1.3. Let $g : (0, 1) \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that

$$f(x, s) = f_0 s^3 + g(x, s)$$

with

$$\lim_{s \rightarrow 0} \frac{g(x, s)}{s^3} = 0 \quad \text{uniformly with respect to all } x \in (0, 1). \quad (4.1)$$

From (4.1), we can see that λg satisfies the assumptions of (1.2). Now, using Theorem 1.1, we have that there are two distinct unbounded continua, \mathcal{C}_k^+ and \mathcal{C}_k^- emanating from $(\mu_k/f_0, 0)$, such that

$$\mathcal{C}_k^v \subset (\{(\mu_k/f_0, 0)\} \cup \Phi_k^v).$$

It is sufficient to show that \mathcal{C}_k^v joins $(\mu_k/f_0, 0)$ to $(\mu_k/f_\infty, \infty)$. Let $(\xi_n, u_n) \in \mathcal{C}_k^v$ where $u_n \not\equiv 0$ satisfies $|\xi_n| + \|u_n\| \rightarrow +\infty$. Proposition 5.1 of [8] implies that $(0, 0)$ is the only solution of (1.5) for $\lambda = 0$, we have $\mathcal{C}_k^v \cap (\{0\} \times X) = \emptyset$. It follows that $\xi_n > 0$ for all $n \in \mathbb{N}$.

Next we show that u_n is one-signed in some interval $(\alpha, \beta) \subseteq (0, 1)$ with $\alpha < \beta$. Let

$$0 < \tau(1, n) < \tau(2, n) < \cdots < \tau(k-1, n) < 1$$

denote the zeros of u_n in $(0, 1)$. Let $\tau(0, n) = 0$ and $\tau(k, n) = 1$. Then, after taking a subsequence if necessary,

$$\lim_{n \rightarrow +\infty} \tau(l, n) = \tau(l, \infty), \quad l \in \{0, 1, \dots, k\}.$$

We claim that there exists $l_0 \in \{0, 1, \dots, k\}$ such that

$$\tau(l_0, \infty) < \tau(l_0 + 1, \infty).$$

Otherwise, we have that

$$1 = \sum_{l=0}^{k-1} (\tau(l+1, n) - \tau(l, n)) \rightarrow \sum_{l=0}^{k-1} (\tau(l+1, \infty) - \tau(l, \infty)) = 0.$$

This is a contradiction. Let $(\alpha, \beta) \subset (\tau(l_0, \infty), \tau(l_0 + 1, \infty))$ with $\alpha < \beta$. For all n sufficiently large, we have $(\alpha, \beta) \subset (\tau(l_0, n), \tau(l_0 + 1, n))$. So u_n does not change its sign in (α, β) .

We claim that there exists a constant M such that $\xi_n \in (0, M]$ for $n \in \mathbb{N}$ large enough. On the contrary, we suppose that $\lim_{n \rightarrow +\infty} \xi_n = +\infty$. Since $(\xi_n, u_n) \in \mathcal{C}_k^v$, it follows that

$$\|u_n\|^2 u_n'' + \xi_n a_n(x) u_n^3 = 0 \quad \text{in } (0, 1),$$

where

$$a_n(x) = \begin{cases} \frac{f(x, u_n)}{u_n^3}, & \text{if } u_n(x) \neq 0, \\ f_0, & \text{if } u_n(x) = 0. \end{cases}$$

From (f1)–(f2), we can see that $\frac{f(x, u_n)}{u_n^3} \geq \sigma$ for some $\sigma > 0$ and all $x \in (0, 1)$, $n \in \mathbb{N}$. So, we have that $\xi_n a_n(x) = +\infty$ for all $x \in (0, 1)$. Applying Theorem 4.1 of [3] on $[\alpha, \beta]$ with $g(x) \equiv \mu_1$, we have that u_n must change its sign in (α, β) for n large enough. This is a contradiction.

Therefore, we get that

$$\|u_n\| \rightarrow +\infty \quad \text{as } n \rightarrow +\infty.$$

Let $h : (0, 1) \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that

$$f(x, s) = f_\infty s^3 + h(x, s)$$

with

$$\lim_{|s| \rightarrow +\infty} \frac{h(x, s)}{s^3} = 0, \quad \lim_{|s| \rightarrow 0} \frac{h(x, s)}{s^3} = f_0 - f_\infty \quad \text{uniformly with respect to all } x \in (0, 1).$$

Then (ξ_n, u_n) satisfies

$$u_n = R_2 (\xi_n f_\infty u_n^3 + h(x, u_n)).$$

Dividing the above equation by $\|u_n\|$ and letting $\bar{u}_n = u_n / \|u_n\|$, we get that

$$\bar{u}_n = R_2 \left(\xi_n f_\infty \bar{u}_n^3 + \frac{h(x, u_n)}{\|u_n\|^3} \right).$$

Let

$$\tilde{h}(x, u) = \max_{0 \leq |s| \leq u} |h(x, s)| \quad \text{for any } x \in (0, 1),$$

then \tilde{h} is nondecreasing with respect to u . Define

$$\bar{h}(x, u) = \max_{u/2 \leq |s| \leq u} |h(x, s)| \quad \text{for any } x \in (0, 1).$$

Then we can see that

$$\lim_{u \rightarrow +\infty} \frac{\bar{h}(x, u)}{u^3} = 0 \quad \text{and} \quad \tilde{h}(x, u) \leq \tilde{h}\left(x, \frac{u}{2}\right) + \bar{h}(x, u).$$

It follows that

$$\limsup_{u \rightarrow +\infty} \frac{\tilde{h}(x, u)}{u^3} \leq \limsup_{u \rightarrow +\infty} \frac{\tilde{h}\left(x, \frac{u}{2}\right)}{u^3} = \limsup_{u/2 \rightarrow +\infty} \frac{\tilde{h}\left(x, \frac{u}{2}\right)}{8\left(\frac{u}{2}\right)^3}.$$

So we have

$$\lim_{u \rightarrow +\infty} \frac{\tilde{h}(x, u)}{u^3} = 0. \quad (4.2)$$

Further it follows from (4.2) that

$$\frac{h(x, u_n)}{\|u_n\|^3} \leq \frac{\tilde{h}(x, |u_n|)}{\|u_n\|^3} \leq \frac{\tilde{h}(x, \|u_n\|_\infty)}{\|u_n\|^3} \leq c^3 \frac{\tilde{h}(x, c\|u_n\|)}{c^3 \|u_n\|^3} \rightarrow 0 \quad \text{as } n \rightarrow +\infty$$

uniformly for $x \in (0, 1)$.

By the compactness of R_2 we obtain that

$$-\|\bar{u}\|^2 \bar{u}'' = \bar{\mu} f_\infty \bar{u}^3,$$

where $\bar{u} = \lim_{n \rightarrow +\infty} \bar{u}_n$ and $\bar{\mu} = \lim_{n \rightarrow +\infty} \bar{\mu}_n$, again choosing a subsequence and relabeling it if necessary. It follows from $\bar{u} = \lim_{n \rightarrow +\infty} \bar{u}_n$ and the triangle inequality that $\|\bar{u}\| = \lim_{n \rightarrow +\infty} \|\bar{u}_n\|$. Since $\|\bar{u}_n\| \equiv 1$, we obtain that $\|\bar{u}\| = 1$. It is clear that $\bar{u} \in \mathcal{C}_k^v$. Theorem 1.2 of [3] shows that $\bar{\mu} = \mu_k / f_\infty$. Therefore, \mathcal{C} joins $(\mu_k / f_0, 0)$ to $(\mu_k / f_\infty, \infty)$. \square

From Theorem 1.3, we can easily get the following corollary.

Corollary 4.1. *Assume that f satisfies (f1)–(f2). Then for*

$$\lambda \in \left(\frac{\mu_k}{f_0}, \frac{\mu_k}{f_\infty} \right) \cup \left(\frac{\mu_k}{f_\infty}, \frac{\mu_k}{f_0} \right),$$

problem (1.5) possesses at least two solutions u_k^+ and u_k^- such that u_k^+ has exactly $k - 1$ simple zeros in $(0, 1)$ and is positive near 0, and u_k^- has exactly $k - 1$ simple zeros in $(0, 1)$ and is negative near 0.

Acknowledgements

The authors wish to express their thanks to the referee for his or her very careful reading of the paper, giving valuable comments and helpful suggestions. The second author is supported by NSF of Liaoning Province (No. 2019-MS-109). The third author is supported by NNSF of China (No. 11871129), the Fundamental Research Funds for the Central Universities (No. DUT20LK04) and Xinghai Youqing funds from Dalian University of Technology.

References

- [1] A. AROSIO, S. PANNIZI, On the well-posedness of the Kirchhoff string, *Trans. Amer. Math. Soc.* **348**(1996), 305–330. <https://doi.org/10.1090/S0002-9947-96-01532-2>; MR1333386
- [2] H. BERESTYCKI, On some nonlinear Sturm–Liouville problems, *J. Differential Equations* **26**(1977), 375–390. [https://doi.org/10.1016/0022-0396\(77\)90086-9](https://doi.org/10.1016/0022-0396(77)90086-9); MR0481230
- [3] X. CAO, G. DAI, Spectrum, global bifurcation and nodal solutions for Kirchhoff type equations, *Electron. J. Differential Equations* **2018**, No. 179, 10 pp. MR3891484
- [4] M. M. CAVALCANTI, V. N. CAVALCANTI, J. A. SORIANO, Global existence and uniform decay rates for the Kirchhoff–Carrier equation with nonlinear dissipation, *Adv. Differential Equations* **6**(2001), 701–730. MR1829093

- [5] C. CHEN, Y. KUO, T. WU, The Nehari manifold for a Kirchhoff type problem involving sign-changing weight functions, *J. Differential Equations* **250**(2011), No. 4, 1876–1908. <https://doi.org/10.1016/j.jde.2010.11.017>; MR2763559
- [6] B. CHENG, X. WU, Existence results of positive solutions of Kirchhoff type problems, *Nonlinear Anal.* **71**(2009), No. 10, 4883–4892. <https://doi.org/10.1016/j.na.2009.03.06>; MR2548720
- [7] G. DAI, Bifurcation and admissible solutions for the Hessian equation, *J. Funct. Anal.* **273**(2017), 3200–3240. <https://doi.org/10.1016/j.jfa.2017.08.001>; MR3695892
- [8] G. DAI, Eigenvalue, global bifurcation and positive solutions for a class of nonlocal elliptic equations, *Topol. Methods Nonlinear Anal.* **48**(2016), 213–233. <https://doi.org/10.12775/TMNA.2016.043>; MR3561429
- [9] G. DAI, Some global results for a class of homogeneous nonlocal eigenvalue problems, *Commun. Contemp. Math.* **21**(2019), No. 3, 1750093, 23 pp. <https://doi.org/10.1142/S0219199717500936>; MR3947059
- [10] G. DAI, Z. FENG, Unilateral global bifurcation for eigenvalue problems with homogeneous operator, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **29**(2019), No. 6, 1950084, 15 pp. <https://doi.org/10.1142/S0218127419500846>; MR3976459
- [11] E. N. DANCER, On the structure of solutions of non-linear eigenvalue problems, *Indiana Univ. Math. J.* **23**(1974), 1069–1076. <https://doi.org/10.1512/iumj.1974.23.23087>; MR0348567
- [12] P. D’ANCONA, S. SPAGNOLO, Global solvability for the degenerate Kirchhoff equation with real analytic data, *Invent. Math.* **108**(1992), 247–262. <https://doi.org/10.1007/BF02100605>; MR1161092
- [13] P. D’ANCONA, Y. SHIBATA, On global solvability of nonlinear viscoelastic equations in the analytic category, *Math. Methods Appl. Sci.* **17**(1994), No. 6, 477–486. <https://doi.org/10.1002/mma.1670170605>; MR1274154
- [14] M. DEL PINO, R. MANÁSEVICH, Global bifurcation from the eigenvalues of the p -Laplacian, *J. Differential Equations* **92**(1991), 226–251. [https://doi.org/10.1016/0022-0396\(91\)90048-E](https://doi.org/10.1016/0022-0396(91)90048-E); MR1120904
- [15] X. HE, W. ZOU, Infinitely many positive solutions for Kirchhoff-type problems, *Nonlinear Anal.* **70**(2009), 1407–1414. <https://doi.org/10.1016/j.na.2008.02.021>; MR2474927
- [16] G. KIRCHHOFF, *Mechanik*, Teubner, Leipzig, 1883.
- [17] Z. LIANG, F. LI, J. SHI, Positive solutions to Kirchhoff type equations with nonlinearity having prescribed asymptotic behavior, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **31**(2014), 155–167. <https://doi.org/10.1016/j.anihpc.2013.01.006>; MR3165283
- [18] Z. LIANG, F. LI, J. SHI, Positive solutions of Kirchhoff type nonlocal elliptic equation: a bifurcation approach, *Proc. Roy. Soc. Edinburgh Sect. A* **147**(2017), No. 4, 875–894. <https://doi.org/10.1017/S0308210516000378>; MR3681567

- [19] J. L. LIONS, On some equations in boundary value problems of mathematical physics, in: *Contemporary developments in continuum mechanics and partial differential equations (Proceedings of the International Symposium on Continuum Mechanics and Partial Differential Equations)*, North-Holland Mathematics Studies, Vol. 30, North-Holland, Amsterdam, 1978, pp. 284–346. [https://doi.org/10.1016/S0304-0208\(08\)70870-3](https://doi.org/10.1016/S0304-0208(08)70870-3)
- [20] T. F. MA, J. E. MUÑOZ RIVERA, Positive solutions for a nonlinear nonlocal elliptic transmission problem, *Appl. Math. Lett.* **16**(2003), No. 2, 243–248. [https://doi.org/10.1016/S0893-9659\(03\)80038-1](https://doi.org/10.1016/S0893-9659(03)80038-1); MR1962322
- [21] A. MAO, Z. ZHANG, Sign-changing and multiple solutions of Kirchhoff type problems without the P.S. condition, *Nonlinear Anal.* **70**(2009), No. 3, 1275–1287. <https://doi.org/10.1016/j.na.2008.02.011>; MR2474918
- [22] K. PERERA, Z. ZHANG, Nontrivial solutions of Kirchhoff-type problems via the Yang index, *J. Differential Equations* **221**(2006), No. 1, 246–255. <https://doi.org/10.1016/j.jde.2005.03.006>; MR2193850
- [23] P. H. RABINOWITZ, Some global results for nonlinear eigenvalue problems, *J. Funct. Anal.* **7**(1971), 487–513. [https://doi.org/10.1016/0022-1236\(71\)90030-9](https://doi.org/10.1016/0022-1236(71)90030-9); MR0301587
- [24] J. SUN, C. TANG, Existence and multiplicity of solutions for Kirchhoff type equations, *Nonlinear Anal.* **74**(2011), No. 4, 1212–1222. <https://doi.org/10.1016/j.na.2010.09.061>; MR2746801
- [25] L. WEN, X. TANG, S. CHEN, Ground state sign-changing solutions for Kirchhoff equations with logarithmic nonlinearity, *Electron. J. Qual. Theory Differ. Equ.* **2019**, No. 47, 1–13. <https://doi.org/10.14232/ejqtde.2019.1.47>; MR3991096
- [26] Z. ZHANG, K. PERERA, Sign changing solutions of Kirchhoff type problems via invariant sets of descent flow, *J. Math. Anal. Appl.* **317**(2006), No. 2, 456–463. <https://doi.org/10.1016/j.jmaa.2005.06.102>; MR2208932